

Multicomponent nonisothermal nucleation. 2. Chapman-Enskog procedure

V.B.Kurasov

Victor.Kurasov@pobox.spbu.ru

This part of the theory directly continues the derivation of the kinetic equation presented in the preprint 990958 of this archive. All definitions are the same and no special remarks are given.

1 Chapman-Enskog procedure

At first we recall the standard version of the Chapman-Enskog procedure and then we shall present the generalization to our situation and fulfill calculations.

Consider equation

$$\frac{\partial P}{\partial t} = -(A + B)P \quad (1)$$

where operators A and B have the following properties

- Operator A depends on variable x , operator B depends on x and y . Both operators don't depend on time t .
- Operator B is supposed to be small in comparison with A . Later we shall rescale time as to have $\|A\| \sim 1$, $\|B\| \ll 1$. The value $\|B\|$ will be the small parameter of the theory.
- The eigenvalues and eigenfunctions of B are unknown. The eigenfunctions A_i and eigenvalues a_i of operator A are supposed to be known. They have the following properties

$$a_0 = 0$$

$$a_i > \delta > 0 \quad i \neq 0$$

with some positive small *delta*. The eigenfunctions are supposed to be normalized.

One can introduce symbol $O(||B||)$ which means that the given expression has the order of $||B||$ or less.

Initial conditions for (1) are supposed to be known. Then one can directly describe the evolution during the first periods of time. Here one can neglect B . One has only to solve equation

$$\frac{\partial P}{\partial t} = -AP \tag{2}$$

Due to the known eigenfunctions of A this solution is quite obvious

$$P = \sum_{m=0}^{\infty} \exp(-a_m t) p_m A_m \tag{3}$$

where

$$p_m = (P|_{t=0}, A_m) \tag{4}$$

will be called the mode amplitudes (A_m will be modes) and the symbol $(,)$ denotes the scalar product.

The final of relaxation according to (3) is obvious

$$P \rightarrow p_0 A_0 \tag{5}$$

The characteristic time of relaxation is

$$t_{rel} = 1/a_1 \tag{6}$$

where a_1 is the smallest eigenvalue of A (except the zero value).

Now $A(p_0 A_0) = 0$ and operator A exhausts its power. One has to consider the action of B ¹.

¹ The physical reason to consider the further stages of evolution is that in the nucleation problem $A_0 = 1$ and the final state of relaxation doesn't provide any benefit into the value of the nucleation rate. It lies in the evident correspondence with the formula $J \sim \exp(F_c)/\Delta\nu$ of the one-dimensional theory where stands the big parameter $\Delta\nu$.

The deviation of P from $p_0 A_0$ can be caused only by the action of operator B . So, in some sense it is small. Then we shall seek the solution in the following form

$$P = p_0 A_0 + \Delta \quad (7)$$

For the correction term Δ in accordance with the perturbation theory one can consider the representation

$$\Delta = \sum_i^{\infty} \delta^{(i)} \quad (8)$$

where every term $\delta^{(i)}$ has the order $\|B\|^i$:

$$\delta^{(i)} \sim O(\|B\|^i)$$

Now we are going to consider the time dependence of the amplitudes p_m . One can note that the action of A leads to the vanishing of p_m for $m \neq 0$. The unique source to fill the given mode is to take into account the action of operator B which is small but still can provide the transition from one mode to another. But B doesn't depend on time. So, the intensities of transitions between modes don't depend on time also. So, the unique source of time dependence is the time dependence of the zero mode p_0 . Later this dependence will be spread by B to all other modes.

As the result we state that the time dependence of an arbitrary mode amplitude is going through the time dependence of p_0 :

$$p_m(t) = p_m(p_0(t)) \quad (9)$$

$$\frac{\partial p_m}{\partial t} = \frac{\partial p_m}{\partial p_0} \frac{\partial p_0}{\partial t} \quad (10)$$

Certainly the time dependence of p_0 isn't fixed and can not be obtained immediately. One has to recalculate it at the every step of approximation². The value $\delta^{(i)}$ depends on time t through $p_0(t)$ in the i -th approximation. At every new³ $\delta^{(i)}$ one has new $p_m^{(i)}$ and new $p_0^{(i)}$. So, one has a new time dependence.

²It is also possible to leave this dependence as some formal parameter and to calculate it when the "final" approximation (i.e. the approximation with the necessary precision) is obtained.

³The lower index denotes modes.

The substitution of expansion

$$P = p_0 A_0 + \sum_{i=1}^{\infty} \delta^{(i)} \quad (11)$$

into (1) gives

$$\frac{\partial p_0}{\partial t} A_0 + \sum_{i=1}^{\infty} \frac{\partial \delta^{(i)}}{\partial t} = -(A + B)(p_0 A_0 + \sum_{i=1}^{\infty} \delta^{(i)}) \quad (12)$$

Note that the i -th term has order i , but the value with the order i will have benefits from all terms with $j \geq i$.

The r.h.s. and l.h.s. of the last equation are some functions and we suppose that they can be decomposed into series over the small parameter $\|B\|$. We shall denote the i -th term which has the order $\|B\|^i$ by the superscript $< i >$. Then one can write for the r.h.s.

$$((A + B)(p_0 A_0 + \sum_{i=1}^{\infty} \delta^{(i)}))^{<i>} = A\delta^{(i)} + B\delta^{(i-1)} \quad (13)$$

The consideration of the l.h.s. is more complicated. At first we have to see the expression for $\partial p_0 / \partial t$. Having projected equation (1) on A_0 one can see that

$$\frac{\partial p_0}{\partial t} = -(A_0, BP) \quad (14)$$

Hence, one can come to the estimate

$$\frac{\partial p_0}{\partial t} = O(\|B\|)$$

One can rewrite the last relation using the decomposition (3)

$$\frac{\partial p_0}{\partial t} = -(A_0, B(p_0 A_0 + \sum_{i=1}^{\infty} \delta^{(i)})) \quad (15)$$

Now it is evident that $\partial p_0 / \partial t$ contains all powers of the small parameter and it has the order $O(\|B\|)$. Now it is worth to determine the j -th term in decomposition over parameter $\|B\|$. It quite easy to do for $\partial p_0 / \partial t$. Namely,

$$(\frac{\partial p_0}{\partial t})^{<i>} = -(A_0, B\delta^{(i-1)}) \quad (16)$$

Then one can fulfill the same transformations for $\partial\delta^{(i)}/\partial t$. One present the following estimate

$$\frac{\partial\delta^{(i)}}{\partial t} = \frac{\partial\delta^{(i)}}{\partial p_0} \frac{\partial p_0}{\partial t} \sim O(\|B\|^i)O(\|B\|) \sim O(\|B\|^{i+1}) \quad (17)$$

So, one can reduce the infinite sum to the finite sum

$$\left(\sum_{j=1}^{\infty} \frac{\partial\delta^{(j)}}{\partial t}\right)^{<i> \rightarrow \left(\sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial t}\right)^{<i> \quad (18)$$

The next transformations are evident

$$\begin{aligned} \left(\sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial t}\right)^{<i> &= \left(\sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial p_0} \frac{\partial p_0}{\partial t}\right)^{<i> = \\ &= \sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial p_0} \left(\frac{\partial p_0}{\partial t}\right)^{<i-j> = \\ &= - \sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial p_0} (A_0, B\delta^{(i-j-1)}) \\ \delta^{(0)} &= p_0 A_0 \end{aligned} \quad (19)$$

Now we can substitute all results into the initial equation and get

$$\begin{aligned} - (A_0, B\delta^{(i-1)}) A_0 - \sum_{j=1}^{i-1} \frac{\partial\delta^{(j)}}{\partial p_0} (A_0, B\delta^{(i-j-1)}) = \\ - A\delta^{(i)} - B\delta^{(i-1)} \end{aligned} \quad (20)$$

As far as the eigenfunctions of A are known one can use the last relation to get $\delta^{(i)}$.

2 Relaxation

Consider now our situation. Certainly, one can choose operator D_1 as the main one.

An operator

$$D_0 = \left(\frac{\partial}{\partial \mu} - 2\mu\right) \frac{\partial}{\partial \mu}$$

has the known set of eigenfunctions which are the Hermite polynomials H_j ($j = 0, 1, 2, 3, 4, \dots$). The scalar product is defined as

$$(\Psi, \Phi) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} dx \exp(-x^2) \Phi(x) \Psi(x)$$

which leads to

$$(H_j, H_i) = \delta_{ij} 2^i i!$$

The eigenvalues of D_0 are

$$\lambda_i = -2i \quad i = 0, 1, 2, \dots$$

One has to mention two important relations

$$\left(\frac{\partial}{\partial \mu} - 2\mu\right) H_i = -H_{i+1}$$

$$\frac{\partial}{\partial \mu} H_i = 2i H_{i-1}$$

The first one gives the way to construct the set of Hermite polynomials. Both these equations allow to introduce in [10] the representation of modes. We shall call the operators in the r.h.s. of two previous equations as the transition operators. The first one is the mode increase operator.

The second one is the mode decrease operator. Then one can consider the operators of multiplication on μ and the differentiation over μ as the superposition of transition operators. Any additional operator appears from the non-Fokker-Planck behavior and from higher terms in the free energy. Non-Fokker-Planck terms give high derivatives. Free energy can be well approximated in the nearcritical region by polynomial and leads to additional multiplications on μ . So, the additional term can be regarded as the superposition of the mode increase operators and mode decrease operators. This procedure resembles the formalism of the secondary quantization. Below we shall use another approach.

It is clear that D_1 can be presented as

$$D_1 = \sum_{l=1}^{\infty} d_l^{(1)} \left(\frac{\partial}{\partial \mu} - 2\mu\right)^l \frac{\partial^l}{\partial \mu^l}$$

with coefficients

$$d_l^{(1)} = - \sum_j \frac{(-\tau_j)^l}{l!} S_{2j} W_j^+ \frac{\tau_j^l}{l!} \quad m = 2, 3, 4, \dots$$

$$\begin{aligned} d_1^{(1)} = & \sum_{j'} W_{j'}^+ \alpha_{acc \ j'} \frac{c_{g \ j'}}{2 \sum_j c_j \nu_j} + \\ & \sum_{i'} W_{i'}^+ (1 - \alpha_{c \ i'}) \alpha_{acc \ i'} \frac{c_{i'}}{2 \sum_j c_j \nu_j} + \\ & \sum_{i'} W_{i'}^+ \alpha_{c \ i'} \frac{c_{i'}}{2 \sum_j c_j \nu_j} - \sum_j (-\tau_j^2) W_j^+ S_{2j} \end{aligned}$$

Then the eigenfunctions of D_1 will be the Hermite polynomials⁴ and the eigenvalues are given by

$$\lambda_j = \sum_{l=1}^j d_l^{(1)} (-2)^l \frac{j!}{(j-l+1)!}$$

As the result it can be seen that one can take D_1 as the main operator in the Chapman-Enskog procedure. The approximate form of kinetic equation will be the following one

$$\frac{\partial P}{\partial t} = D_1 P$$

But already in the investigation of the stationary distribution establishing one will see that the formal priority of D_1 in comparison with D_2 isn't sufficient to ensure the relaxation to the stationary state.

Operator D_2 will be small in comparison with D_1 in terms of some small parameter (not in formal sense described above) only when τ_j far all j are small parameters. Then the main terms in operator D_2 will be the terms with the smallest sum of indexes $l + m$. Then the required condition of the smallness of D_2 formulated in terms of the relaxation times will be

$$p_{rel} \equiv \frac{|-\sum_j \tau_j^3 W_j^+|}{|\lambda_1|} \ll 1$$

⁴In the main order S_{2j} goes to 1.

This parameter has been required⁵ to be small in [8]. Then the solution at the relaxation stage can be written as the series of the relaxation modes

$$P = \sum_{l=0}^{\infty} p_l^{(0)} \exp(-\lambda_l t) H_l$$

where

$$p_l^{(0)} = (2^l l!)^{-1} (H_l, P(t=0))$$

are initial amplitudes. This series ensures the relaxation to the stationary state $p_0^{(0)} H_0$:

$$P \rightarrow p_0^{(0)} H_0 \equiv P_{rel}$$

The time of relaxation t_{rel} is given by

$$t_{rel} \approx \lambda_1^{-1}$$

The mentioned restriction $p_{rel} \leq 1$ (actually according to [8] one has to require $p_{rel} \ll 1$) practically excludes the nucleation under the strong thermal effects. So, the situation considered in [8] is rather poor.

One can note that the final state P_{rel} corresponds to the equilibrium distribution over μ . So, it is the eigenfunction of the operator $D_1 + D_2$ with the zero eigenvalue

$$(D_1 + D_2)P_{rel} = 0$$

This can be easily proven when we return to the form of the finite differences and reconstruct D_1 and D_2 .

Certainly it is difficult to determine all eigenfunctions and eigenvalues of $D_1 + D_2$. We needn't all relaxation modes but only the final one. Also we have to estimate the time of relaxation. One can prove that the relaxation time $t_{rel} \approx \lambda_1^{-1}$ for equation

$$\frac{\partial P}{\partial t} = (D_1 + D_2)P$$

is less or equal to the previous relaxation time $t_{rel} \approx \lambda_1^{-1}$

$$t_{rel} \approx \lambda_1^{-1} \leq t_{rel} \approx \lambda_1^{-1}$$

⁵with account of the mentioned error.

Operator D_2 "helps" the relaxation to P_{rel} . The way to prove the last estimate is to consider the blocks along μ axis and estimate the action of D_2 between blocks.

As far as D_3 , D_4 have small parameter in comparison with D_1 then no special condition is required. Now we can consider $D_1 + D_2$ as the main operator at the relaxation stage. The smallness of p_{rel} isn't now required.

3 Correction terms

When P is close to P_{rel} which is the eigenfunction with the zero eigenvalue the operators D_1 or $D_1 + D_2$ can not be considered as the main ones. One has to use the Chapman-Enskog asymptotic decomposition. But to fulfill this decomposition one has to know all eigenfunctions of the main operator. So we can not use $D_1 + D_2$ as the main operator now.

We shall redefine the main operator after the end of the relaxation stage. Now the main operator will be D_1 . One can see that "correction operator" $D_2 + D_3 + D_4$ has a complicate structure. Operators D_3 and D_4 have small parameter. Operator D_2 is small only in the formal sense. So, the natural modification of the Chapman-Enskog procedure is to consider two Chapman-Enskog procedures. One can include the whole internal Chapman-Enskog procedure into every step of the external Chapman-Enskog procedure. The initial procedure has the aim to "invert" the action of operator D_2 and the external procedure will "invert" the action of $D_3 + D_4$.

Every new approximation will contain the small parameter $1/\Delta\kappa \sim \kappa^{2/3}$ (or at least $\Delta\kappa/\kappa \sim \kappa^{-1/3}$) in the order of the number of external approximation. This order decrease the order of flow calculated in the initial (zero) approximation. But one has to stop the external procedure until the resulting order is greater than $\kappa^0 = 1$ because the continuous description along ν_i will be violated in this order. One has to note that the evolution along ν_i is principally different in comparison with evolution along μ . The domain along ν_i is concentrated only at the integer values. The value of μ is the real value. This difference leads to the specific lattice corrections described in [10]. These corrections have very complicated structure, an account can not be fulfilled explicitly in the analytic way. As the result we have to restrict ourselves by the first correction and adopt that the account of the first correction in the external procedure is already sufficient. But in the internal

procedure we have to calculate the great number of approximations.

Also one has to note here the requirement to have no operators S_{1j} , $S_{@j}$, S_{3j} in the final results. Really, the presence of this operators corresponds to the elementary shift taken into account. This lies in contradiction with the lattice domain along ν_i .

The matter under discussion is the rate of nucleation and to determine this rate and need only the embryos flows averaged over μ . This operation corresponds to the projection on $H_0 \sim const$.

Now we can turn ourselves to establish equations for $\partial p_0 / \partial t$. The last dependence is extremely important because in the Chapman-Enskog procedure all dependence of approximations on time is going through dependence of p_0 on time. To get $\partial p_0 / \partial t$ one can project kinetic equation

$$\frac{\partial P}{\partial t} = [D_1 + D_2 + D_3 + D_4]P$$

on H_0 i.e. fulfill the integration $\int_{-\infty}^{\infty} \exp(-\mu^2) \dots d\mu$. The l.h.s. gives $\partial p_0 / \partial t$. Then

$$\frac{\partial p_0}{\partial t} = (H_0, [D_1 + D_2 + D_3 + D_4]P)$$

The r.h.s. will lead to more complicate expression. Consider

$$(H_0, D_1 P) = (H_0, D_1 \sum_i p_i H_i) = (H_0, \sum_i \lambda_i p_i H_i) = \lambda_0 p_0 = 0$$

as far as $\lambda_0 = 0$. Consider

$$(H_0, D_2 P) = (H_0, D_2 \sum_{i=0}^{\infty} p_i H_i)$$

From explicit form of D_2 one can see that the last elementary operator in action of D_2 will be the mode increase operator $(\frac{\partial}{\partial \mu} - 2\mu)$. So, if we present $D_2 \sum_{i=0}^{\infty} p_i H_i$ as

$$D_2 \sum_{i=0}^{\infty} p_i H_i = \sum_{i=0}^{\infty} c_i H_i$$

then $c_0 = 0$. There is no projection on the zero mode. Then

$$(H_0, \sum_{i=1}^{\infty} c_i H_i) = 0$$

for arbitrary c_i . Then

$$(H_0, D_2 P) = 0$$

Consider $(H_0, D_3 P)$. Operator D_3 can be split into two parts D_{3a} and D_{3b} defined as

$$\begin{aligned} D_{3a} &= \sum_j \sum_{l=1}^{\infty} \frac{(-\tau_j)^l}{l!} \left(\frac{\partial}{\partial \mu} - 2\mu \right)^l L_j S_{1j} P(\{\nu_i\}, \mu) \\ D_{3b} &= - \sum_j \frac{\partial}{\partial \nu_j} W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m} P(\{\nu_i\}, \mu) \\ D_3 &= D_{3a} + D_{3b} \end{aligned}$$

Then

$$(H_0, D_{3a} P) = 0$$

and the reasons are the same as for D_2 . The action of D_{3b} can be presented as

$$(H_0, D_{3b} P) = - \sum_j \frac{\partial}{\partial \nu_j} S_{3j} (H_0, W_j^+ \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m} P)$$

The last operator D_4 gives

$$(H_0, D_4 P) = - \sum_j \frac{\partial}{\partial \nu_j} (H_0, L_j P)$$

As the result one can get

$$\frac{\partial p_0}{\partial t} = - \sum_j \frac{\partial}{\partial \nu_j} (H_0, [L_j + W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m}] P)$$

The values in the r.h.s. of the last equation can be interpreted as the flows J_j along ν_j axis

$$J_j = (H_0, [L_j + W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m}] P)$$

which transfers the last equation in the standard form

$$\frac{\partial p_0}{\partial t} = - \sum_j \frac{\partial}{\partial \nu_j} J_j \tag{21}$$

4 Calculations

Now we can turn to the direct calculations. The distribution P can be presented into the following form

$$P = p_0 H_0 + \sum_{l=0}^{\infty} p_l(p_0) H_l$$

$$p_l(p_0) = \sum_{m=1}^{\infty} p_l^{(m)}(p_0)$$

where the lower index indicates the number of mode and the upper index in brackets indicates the number of approximation. As far as we already stated we can fulfill in the external procedure only one step and it isn't necessary to mark it. The upper index, thus, corresponds to the internal approximations.

Now we can rewrite expression for J_j in terms of announced decomposition. Then

$$J_j = (H_0, [L_j + W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m}] [p_0 H_0 + \sum_{l=0}^{\infty} p_l(p_0) H_l])$$

Consider various terms of the last expression. Namely,

$$(H_0, L_j P) = L_j p_0$$

Then consider $(H_0, W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m} [p_0 H_0 + \sum_{l=0}^{\infty} p_l(p_0) H_l])$ The term $p_0 H_0$ doesn't leads to any influence because in $\sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m}$ there is at least one operator of the mode decrease. Then

$$\begin{aligned} (H_0, W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m} [p_0 H_0 + \sum_{l=0}^{\infty} p_l(p_0) H_l]) &= (H_0, W_j^+ S_{3j} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial^m}{\partial \mu^m} \sum_{l=0}^{\infty} p_l(p_0) H_l) = \\ &= (H_0, W_j^+ S_{3j} \sum_{i=1}^{\infty} \sum_{m=1}^{\infty} \frac{\tau_j^m}{m!} \frac{\partial}{\partial \mu^m} p_i(p_0) H_i) = W_j^+ S_{3j} \sum_{m=1}^{\infty} \tau_j^m p_m(p_0) 2^m \end{aligned}$$

The resulting expression for J_j will be the following

$$J_j = L_j p_0 + W_j^+ S_{3j} \sum_{m=1}^{\infty} \tau_j^m p_m(p_0) 2^m$$

In the main order we can omit S_{3j} . Certainly, the correction term in S_{3j} has the order greater than L_j has but there is no zero mode except $L_j p_0$. So, we have to keep L_j and throw away S_{3j} here. Then

$$J_j = L_j p_0 + W_j^+ \sum_{m=1}^{\infty} \tau_j^m p_m(p_0) 2^m$$

Now we can calculate the set of corrections. The initial condition for the asymptotic expansion will be the result of the relaxation stage, i.e. P_{rel} . One can easily note that

$$D_1 P_{rel} = 0$$

$$D_2 P_{rel} = 0$$

$$D_{3b} P_{rel} = 0$$

because the first elementary operator is the mode decrease operator. In D_{3a} there are only mode increase operators and then

$$\begin{aligned} D_{3a} P_{rel} &= \sum_j L_j S_{1j} \sum_{l=1}^{\infty} \frac{(-\tau_j)^l}{l!} \left(\frac{\partial}{\partial \mu} - 2\mu \right)^l p_0 H_0 = \\ &= \sum_j L_j S_{1j} \sum_{l=1}^{\infty} \frac{(-\tau_j)^l}{l!} (-1)^l p_0 H_l = \sum_j L_j S_{1j} \sum_{l=1}^{\infty} \frac{(\tau_j)^l}{l!} p_0 H_l \end{aligned}$$

The last operator contains the small parameter in the first order. In the main order one can throw S_{1j} out.

Operator D_4 has at least the small parameter r in power 2. So, it is small in comparison with D_{3a} which has r in the first power. As far as operators D_1 , D_2 have no action we have to take into account D_3 but we can neglect the action of D_4 .

As the result we have

$$p_l^{(1)} = \frac{1}{\lambda_l l!} \sum_j \tau_j^l L_j p_0$$

In the zero approximation

$$\left(\frac{\partial p_0}{\partial t} \right)^{(0)} = 0$$

The structure of first approximation is rather simple due to the zero approximation which is localized only at the zero mode. Now the current approximation has all modes and operator D_2 will lead to the non-zero result. So, the operator D_2 will be the main one and one can neglect D_3 and D_4 . In D_2 , thus, one can neglect S_{2j}

The time derivative can be also neglected. Really, due to (15)

$$\left(\frac{\partial p_0}{\partial t}\right)^{(1)} \sim -(H_0, [D_2 + D_3 + D_4]H_0)$$

in the main order.

One can easily see that

$$(H_0, D_2 H_0) = 0$$

$$(H_0, D_{3b} H_0) = 0$$

as far as the first operator in the action of D_2 and D_{3b} is the derivative over μ which gives zero in application to $H_0 \sim \text{const}$.

As far D_{3a} has the small parameter and D_4 has the result $(\partial p_0 / \partial t)^{(1)}$ is small. As far as we calculate in the main order of small parameter we can neglect $(\partial p_0 / \partial t)^{(1)}$.

In all approximations from the second one we can use (21) and see due to the smallness of $\partial / \partial \nu_i$ the smallness of $(\partial p_0 / \partial t)^{(i)}$ $i = 2, 3, 4, \dots$

We see that the Chapman-Enskog procedure is now reduced to the trivial equation (except the zero mode)

$$p^{(i+1)} = D_1^{-1} D_2 p^{(i)}$$

with the evident initial approximation. In the main order one can take S_{2j} away from the operator D_2 . This form can be obtained from the simple analysis without Chapman-Enskog formalism.

We can present the last equation in the following form

$$p_k^{(i+1)} = \sum_{q=1}^{\infty} \Gamma_{kq} p_q^{(i)}$$

as the matrix representation of the linear operator $D_1^{-1} D_2$ in the basis H_i (without the zero mode). All Γ_{qq} can be put to zero.

One can note the following important features: 1). Any operator acts until the current moment in the already fulfilled part of the external Chapman-Enskog procedure only one time. 2). Every operator has its own specific

structure and can not be reproduced by the actions of other operators. Both these features allow to forget about S_{1j} , S_{2j} , S_{3j} . If any operator had been used more than one time then one would take these operators into account.

The absence of S_{1j} , S_{2j} , S_{3j} is very important in the context of the lattice structure of the domain in the $\{\nu_i\}$ plane. Only the absence of these operators allows to ignore this lattice structure.

5 Final expressions

Now we have to establish the expression for Γ_{kq} . As far as D_2 is the superposition of the mode decrease operators and mode increase operators it will transfer H_i into superposition of H_i . The action of D_1^{-1} on H_j (except zero mode) is evident - it is multiplication H_j on λ_j^{-1} .

How D_2 transfers H_q into H_k ? At first according to the definition of D_2 the mode decrease operators will act. The number l of the mode decrease operators will be between l_0 and q . Parameter l_0 appeared from the evident requirement that we have to fall lower than the k -th level. So, $l_0 = 1$ if $k > q$ and $l_0 = q - k + 1$ if $q > k$. The result of the action of the mode decrease operators will give the coefficient $2^l q! / (q - l)!$. Then to get H_k one has to apply $k - (q - l)$ mode increase operators which will give coefficient $(-1)^{k-(q-l)}$. As the result

$$\Gamma_{kq} = \sum_j W_j^+ \sum_{l=l_0}^q \frac{\tau_j^{k-q+l}}{(k-q+l)!} \frac{\tau_j^l}{l!} 2^l \frac{q!}{(q-l)!}$$

Also one can consider another representation of Γ_{kq} . At first we fall from the q level to the l level which gives $\frac{\tau_j^{q-l}}{(q-l)!} \frac{q!}{l!} 2^{q-l}$. then we increase the mode from the l level to the k level which gives $\frac{\tau_j^{k-l}}{(k-l)!}$. The result will be

$$\Gamma_{kq} = \sum_j W_j^+ \sum_{l=0}^{\min\{q,k\}=1} \frac{\tau_j^{q-l}}{(q-l)!} \frac{q!}{l!} 2^{q-l} \frac{\tau_j^{k-l}}{(k-l)!}$$

Note that Γ_{kq} doesn't depend on the number of approximation. This will lead to some important consequences. The first one is the possibility to write equation between p_k :

$$p_k = \sum_{q=1}^{\infty} \Gamma_{kq} p_q + p_k^{(1)}$$

or

$$p_k = \sum_{q=1}^{\infty} \Gamma_{kq} p_q + \frac{1}{\lambda_k k!} \sum_j \tau_j^k L_j p_0$$

Due to the linearity of equations the decomposition of the initial approximation will be reproduced in all approximations. Namely, we shall present $p_l^{(1)}$ in the following form

$$p_l^{(1)} = \sum_j p_{lj}^{(1)}$$

where

$$p_{lj}^{(1)} = \frac{1}{\lambda_l l!} \tau_j^l L_j p_0$$

and the second lower index indicates component in the initial approximation (later Γ_{kq} will mixture different components).

The linearity results in the possibility of decomposition

$$p_k^{(i)} = \sum_j p_{kj}^{(i)}$$

Index j has no correspondence here with the direct decomposition over components.

For $p_{kj}^{(i)}$ the following recurrent expression

$$p_{kj}^{(i+1)} = \sum_{q=1}^{\infty} \Gamma_{kq} p_{qj}^{(i)}$$

is valid.

The total amplitude p_k can be also decomposed as

$$p_k = \sum_j p_{kj}$$

where p_{kj} are

$$p_{kj} = \sum_{i=1}^{\infty} p_{kj}^{(i)}$$

For p_{kj} one can write

$$p_{kj} = \sum_{q=1}^{\infty} \Gamma_{kq} p_{qj} + p_{kj}^{(1)}$$

One can also move operators L_j through $\hat{\Gamma} \equiv \{\Gamma_{kq}\}$. This gives the series of relations. For the first approximation one can write

$$p_l^{(1)} = \sum_j L_j a_{lj}^{(1)} p_0$$

where

$$a_{lj}^{(1)} = \tau_j^l \frac{1}{\lambda_l l!}$$

The recurrent relations for $a_{lj}^{(i)}$ will be

$$a_{kj}^{(i+1)} = \sum_{q=1}^{\infty} \Gamma_{kq} a_{qj}^{(i)}$$

In terms of $a_{kj}^{(i)}$ the value $p_k^{(i)}$ can be easily expressed as

$$p_k^{(i)} = \sum_j a_{kj}^{(i)} L_j p_0$$

Having introduced

$$a_{kj} = \sum_{i=1}^{\infty} a_{qj}^{(i)}$$

one can get

$$a_{kj} = \sum_{q=1}^{\infty} \Gamma_{kq} a_{qj} + a_{kj}^{(1)}$$

In terms of a_{kj} the value p_k can be easily expressed as

$$p_k = \sum_j a_{kj} L_j p_0$$

The given decompositions are rather attractive from the first point of view, but actually one can not simplify the problem with the help of these decompositions.

The equation for p_0 will be the following one

$$\frac{\partial p_0}{\partial t} = - \sum_j \frac{\partial}{\partial \nu_j} \sum_i (\delta_{ij} + W_j^+ S_{3j} \sum_{m=1}^{\infty} \tau_j^m 2^m a_{mi}) L_i p_0$$

and in the main order

$$\frac{\partial p_0}{\partial t} = - \sum_j \frac{\partial}{\partial \nu_j} \sum_i (\delta_{ij} + W_j^+ \sum_{m=1}^{\infty} \tau_j^m 2^m a_{mi}) L_i p_0$$

where indexes i and j marks components. The last equation has the standard form investigated in [6]. The part of [8] concerning the solution of the last equation in the two dimensional sense is also acceptable as far as it reproduces [11] even in details⁶.

Now we shall present another method to calculate correction terms in the Chapman-Enskog procedure.

One can easily note that recurrent equations for $p_l^{(i)}$ and for $p_l^{(i)}{}_j$ will lead to

$$\begin{aligned} \lim_{i \rightarrow \infty} p_l^{(i)}{}_j &\rightarrow \text{const}_l{}_j (\gamma_{max})^i \\ \lim_{i \rightarrow \infty} p_l^{(i)} &\rightarrow \text{const}_l (\gamma_{max})^i \end{aligned}$$

where γ_{max} is the eigenvalue of $\hat{\Gamma} \equiv \{\Gamma_{pq}\}$ with the maximal absolute value.

Then one can say that starting from some number m_{lim} the tails of sums $\sum_i p_l^{(i)}{}_j$, $\sum_i p_l^{(i)}$ resemble the tail of geometric progressions. Then one can easily calculate these sums

$$\sum_{i=m_{lim}}^{\infty} p_l^{(i)}{}_j \sim \frac{p_l^{(m_{lim})}{}_j}{1 - \gamma_{max}}$$

and

$$\sum_{i=m_{lim}}^{\infty} p_l^{(i)} \sim \frac{p_l^{(m_{lim})}}{1 - \gamma_{max}}$$

The first m_{lim} terms have to be calculated explicitly. The boundary m_{lim} which depends on j, l can be found as the characteristic boundary when $\hat{\Gamma}^{i+1}/\hat{\Gamma}^i$ approaches some constant independent on i . This procedure gives also the value of γ_{max} .

Certainly, the most interesting situation is the strong manifestation of the thermal effects. Here the thermal effects can not be considered as some corrections but radically change the character of the process.

⁶except several misprints appeared in [8]. Also the initial matrix (with dimension 2) isn't diagonal which doesn't produce any difficulties.

Really, the main dependence of the nucleation rate is accumulated in exponent of the critical embryo free energy. The giant renormalization due to the thermal effects means that the temperature of the critical embryo really differs from the temperature of the vapor-gas media. The relative difference of the temperature expressed in the units of β (in estimates we can forget about different components) can attain several units. It means that the characteristic value μ_0 of μ attains several units.

One can easily note that to reproduce the real solution (i.e. to attain the values of μ in several units) we have to take into account a great number of modes. Really, the Hermite polynomials H_n are the polynomials of power n . The characteristic region of localization of the function $H_n(\mu) \exp(-\mu^2)$ is $[\sim -\sqrt{n}, \sim \sqrt{n}]$. So, to describe the situation correctly we have to take into account at least μ_0^2 modes. This quantity equals to the dimension of the matrix $\hat{\Gamma}$ which becomes also great. This produces numerical difficulties.

Consider the matrix $\hat{\Gamma}$. Due to factorials in the denominators the limit of the elements with a big indexes is going to zero. It takes place under the arbitrary τ_j . The r.h.s. of the matrix equation, i.e. the known vector $p^{(1)} = \{p_i^{(0)}\}$ also has vanishing elements when $i \rightarrow \infty$. The structure of the matrix $\hat{\Gamma}$ is the following: the upper triangle matrix has the elements with big values due to the big coefficient appeared from the action of the mode decrease. The lower triangle matrix is "smaller" than the upper one. Thus, the following method will be rather effective⁷: one can split matrix $\hat{\Gamma}$ into the upper triangle matrix (with big elements) and the lower triangle matrix (with moderate elements). Then at every step of the iteration procedure the upper triangle matrix will be inverted (it is easy to do). The zero mode (in fact it is the first line in the matrix equation) will be calculated explicitly.

One can easily note the the iteration solutions of the matrix equation return us to the initial formulation of the problem at the level of the recurrent relations. So, the presented approach to calculate the maximal eigenvalue gets now a solid ground. It is also more effective because it allows to estimate the necessary number of modes taken into account directly (by the smallness of the higher mode at every step).

⁷With the special account of the Chapman-Enskog specific features

References

- [1] A. Kantrowitz, J. Chem. Phys. 19 (1951) 1097.
- [2] Kuni F.M., preprint, Institute for Theoretical physics, Kiev, ITP-83-80R, 1983.
- [3] J. Feder, K.C. Russel, J.Lothe, G.M. Pound, Adv. Phys. 15 (1996) 111.
- [4] A.P. Grinin, F.M. Kuni, Theor. Mat. Phys. 80 (1989) 968.
- [5] J.S. Langer, Ann. Phys. 54 (1969) 258.
- [6] F.M. Kuni, A.A. Melikhov, Theor. Mat. Phys. 83 (1990) 530.
- [7] M. Lazaridis, Y. Drossinos, J. Phys. A 30 (1997) 3847.
- [8] Y.S.Dzhikaiev, J. Teichman, M. Grmela, Physica A 267 (1999) 322.
- [9] F.M. Kuni, A.P. Grinin, A.K. Shchekin, Physica A 252 (1998) 67.
- [10] V.B. Kurasov Universality in kinetics of the first order phase transitions, Chemistry Research Institute of St. Petersburg University, St.Petersburg, 1997, 400 p.
- [11] A.A. Melikhov, V.B. Kurasov, Yu. Sh. Dzhikaiev, F.M. Kuni, Sov. Phys. Techn. Phys. 36 (1991) 14.
- [12] R.Strey, Y. Viisanen, J. Chem. Phys. 99 (1993) 4693.
- [13] D. Stauffer, J. Aerosol Sci. 7 (1976) 319.
- [14] H. Reiss, J. Chem.Phys. 18 (1950) 840.